

Answers To Calculus Of A Single Variable

1 □ FUNCTIONS AND MODELS

1.1 Four Ways to Represent a Function

1. The functions $f(x) = x + \sqrt{2-x}$ and $g(u) = u + \sqrt{2-u}$ give exactly the same output values for every input value, so f and g are equal.
2. $f(x) = \frac{x^3 - x}{x - 1} = \frac{x(x-1)}{x-1} = x$ for $x - 1 \neq 0$, so f and g (where $g(x) = x$) are not equal because $f(1)$ is undefined and $g(1) = 1$.
3. (a) The point $(-2, 2)$ lies on the graph of g , so $g(-2) = 2$. Similarly, $g(0) = -2$, $g(2) = 1$, and $g(3) \approx 2.5$.
(b) Only the point $(-4, 3)$ on the graph has a y -value of 3, so the only value of x for which $g(x) = 3$ is -4 .
(c) The function outputs $g(x)$ are never greater than 3, so $g(x) \leq 3$ for the entire domain of the function. Thus, $g(x) \leq 3$ for $-4 \leq x \leq 4$ (or, equivalently, on the interval $[-4, 4]$).
(d) The domain consists of all x -values on the graph of g : $\{x \mid -4 \leq x \leq 4\} = [-4, 4]$. The range of g consists of all the y -values on the graph of g : $\{y \mid -2 \leq y \leq 3\} = [-2, 3]$.
(e) For any $x_1 < x_2$ in the interval $[0, 2]$, we have $g(x_1) < g(x_2)$. [The graph rises from $(0, -2)$ to $(2, 1)$.] Thus, $g(x)$ is increasing on $[0, 2]$.
4. (a) From the graph, we have $f(-4) = -2$ and $g(3) = 4$.
(b) Since $f(-3) = -1$ and $g(-3) = 2$, or by observing that the graph of g is above the graph of f at $x = -3$, $g(-3)$ is larger than $f(-3)$.
(c) The graphs of f and g intersect at $x = -2$ and $x = 2$, so $f(x) = g(x)$ at these two values of x .
(d) The graph of f lies below or on the graph of g for $-4 \leq x \leq -2$ and for $2 \leq x \leq 3$. Thus, the intervals on which $f(x) \leq g(x)$ are $[-4, -2]$ and $[2, 3]$.
(e) $f(x) = -1$ is equivalent to $y = -1$, and the points on the graph of f with y -values of -1 are $(-3, -1)$ and $(4, -1)$, so the solution of the equation $f(x) = -1$ is $x = -3$ or $x = 4$.
(f) For any $x_1 < x_2$ in the interval $[-4, 0]$, we have $g(x_1) > g(x_2)$. Thus, $g(x)$ is decreasing on $[-4, 0]$.
(g) The domain of f is $\{x \mid -4 \leq x \leq 4\} = [-4, 4]$. The range of f is $\{y \mid -2 \leq y \leq 3\} = [-2, 3]$.
(h) The domain of g is $\{x \mid -4 \leq x \leq 3\} = [-4, 3]$. Estimating the lowest point of the graph of g as having coordinates $(0, 0.5)$, the range of g is approximately $\{y \mid 0.5 \leq y \leq 4\} = [0.5, 4]$.
5. From Figure 1 in the text, the lowest point occurs at about $(t, a) = (12, -85)$. The highest point occurs at about $(17, 115)$. Thus, the range of the vertical ground acceleration is $-85 \leq a \leq 115$. Written in interval notation, the range is $[-85, 115]$.

Answers to calculus of a single variable can be both enlightening and challenging for students and enthusiasts alike. Calculus, particularly single-variable calculus, serves as a foundation for numerous scientific and engineering disciplines. This branch of mathematics deals with functions, limits, derivatives, integrals, and infinite series, all of which are essential for understanding change and motion. In this article, we will explore the fundamental concepts of single-variable calculus, provide solutions to common problems, and discuss applications that highlight the importance of this mathematical tool.

Understanding the Basics of Single Variable Calculus

Single-variable calculus focuses on functions of one variable. It is primarily concerned with two main concepts: differentiation and integration.

1. Differentiation

Differentiation involves finding the derivative of a function, which represents the rate of change of the function with respect to its variable. The derivative is denoted as $f'(x)$ or $\frac{df}{dx}$.

Key Concepts of Differentiation:

- Definition of Derivative: The derivative of a function $f(x)$ at a point x is defined as:

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

- Rules of Differentiation:

- Power Rule: If $f(x) = x^n$, then $f'(x) = n x^{n-1}$.

- Product Rule: If $f(x) = u(x) \cdot v(x)$, then $f'(x) = u'(x) v(x) + u(x) v'(x)$.

- Quotient Rule: If $f(x) = \frac{u(x)}{v(x)}$, then $f'(x) = \frac{u'(x)v(x) - u(x)v'(x)}{(v(x))^2}$.

- Chain Rule: If $f(x) = g(h(x))$, then $f'(x) = g'(h(x)) \cdot h'(x)$.

2. Integration

Integration is the reverse process of differentiation. It involves finding the integral of a function, which represents the accumulation of quantities and the area under a curve.

Key Concepts of Integration:

- Definite Integral: The definite integral of $f(x)$ from a to b is given by:

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

where $F(x)$ is an antiderivative of $f(x)$.

- Indefinite Integral: The indefinite integral is represented as:

$$\int f(x) \, dx = F(x) + C$$

where C is a constant of integration.

Integration Techniques:

- Substitution Method: Used to simplify integrals by changing variables.
- Integration by Parts: Based on the product rule for differentiation, it is defined as:

$$\int u \, dv = uv - \int v \, du$$

- Partial Fraction Decomposition: Used for integrating rational functions by expressing them as a sum of simpler fractions.

Common Problems and Their Solutions

Understanding the basic concepts is essential, but applying them to solve problems is where mastery lies. Here are some common types of problems encountered in single-variable calculus along with their solutions.

1. Finding Derivatives

Problem: Find the derivative of the function $f(x) = 3x^4 - 5x^3 + 2x - 7$.

Solution:

Using the power rule:

$$f'(x) = 12x^3 - 15x^2 + 2$$

2. Evaluating Definite Integrals

Problem: Evaluate the integral $\int_1^3 (2x^2 + 3) \, dx$.

Solution:

1. Find the antiderivative:

$$\int (2x^2 + 3) \, dx = \frac{2}{3}x^3 + 3x + C$$

2. Apply the limits:

$$\begin{aligned} & \left[\frac{2}{3}(3^3) + 3(3) \right] - \left[\frac{2}{3}(1^3) + 3(1) \right] = \\ & \left[\frac{2}{3} \cdot 27 + 9 \right] - \left[\frac{2}{3} \cdot 1 + 3 \right] = \\ & \left[18 + 9 \right] - \left[\frac{2}{3} + 3 \right] = 27 - \left[\frac{2}{3} + \frac{9}{3} \right] = 27 - \frac{11}{3} = \frac{81}{3} - \frac{11}{3} = \frac{70}{3} \end{aligned}$$

\]

3. Solving Rates of Change Problems

Problem: A balloon is being inflated, and its radius increases at a rate of 2 cm/min. Find the rate of change of the volume of the balloon when the radius is 5 cm.

Solution:

1. The volume (V) of a sphere is given by:

$$V = \frac{4}{3}\pi r^3$$

\]

2. Differentiate with respect to time (t) :

\[

$$\frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt}$$

\]

3. Substitute $(r = 5)$ cm and $(\frac{dr}{dt} = 2)$ cm/min:

\[

$$\frac{dV}{dt} = 4\pi (5^2)(2) = 4\pi (25)(2) = 200\pi \text{ cm}^3/\text{min}$$

\]

Applications of Single Variable Calculus

Single-variable calculus is not just an academic exercise; it has vast applications in various fields.

1. Physics

In physics, calculus is used to describe motion. For example:

- Velocity: The derivative of the position function gives the velocity function.
- Acceleration: The derivative of the velocity function gives the acceleration function.

2. Economics

In economics, calculus is utilized in various ways:

- Marginal Cost and Revenue: The derivative of the cost function gives the marginal cost, while the derivative of the revenue function gives the marginal revenue.
- Optimization: Calculus helps in finding the maximum profit or minimum cost by analyzing the critical points of profit and cost functions.

3. Biology

In biology, calculus is applied in population modeling:

- Exponential Growth Models: Differential equations can describe how populations grow over time.
- Rate of Change in Reactions: Calculus is used in pharmacokinetics to model the rate at which drugs enter and leave the body.

Conclusion

Answers to calculus of a single variable encompass a vast array of concepts, techniques, and applications. From understanding derivatives and integrals to solving real-world problems, calculus provides the necessary tools to analyze and interpret change. Mastering these concepts not only enhances mathematical skills but also opens doors to various fields of study and professional opportunities. Whether you are a student preparing for exams or an enthusiast seeking to deepen your understanding, embracing the beauty of single-variable calculus will undoubtedly enrich your intellectual journey.

Frequently Asked Questions

What is the fundamental theorem of calculus?

The fundamental theorem of calculus links the concept of differentiation and integration, stating that if F is an antiderivative of f on an interval $[a, b]$, then the integral of f from a to b is equal to $F(b) - F(a)$.

How can I find the derivative of a function using the limit definition?

The derivative of a function f at a point x is defined as the limit: $f'(x) = \lim_{h \rightarrow 0} [(f(x + h) - f(x)) / h]$.

What are the common rules for differentiation?

Common rules for differentiation include the power rule, product rule, quotient rule, and chain rule, which help simplify the process of finding derivatives.

What is the difference between definite and indefinite integrals?

An indefinite integral represents a family of functions and includes a constant of integration, while a definite integral calculates the net area under a curve over a specific interval $[a, b]$.

How do I evaluate limits that result in indeterminate forms?

To evaluate limits that result in indeterminate forms like $0/0$ or ∞/∞ , you can use L'Hôpital's rule, factorization, or algebraic simplification to resolve the indeterminacy.

What is the purpose of the mean value theorem?

The mean value theorem states that if a function is continuous on a closed interval $[a, b]$ and differentiable on the open interval (a, b) , then there exists at least one c in (a, b) such that $f'(c) = (f(b) - f(a)) / (b - a)$.

How do you apply integration techniques like substitution and integration by parts?

Substitution helps simplify integrals by changing variables, while integration by parts, derived from the product rule, is used to integrate products of functions and is expressed as $\int u \, dv = uv - \int v \, du$.

Find other PDF article:

<https://soc.up.edu.ph/09-draft/Book?ID=nHw78-4813&title=bible-study-on-relationship-with-god.pdf>

Answers To Calculus Of A Single Variable

Answers - The Most Trusted Place for Answering Life's Questions

Answers is the place to go to get the answers you need and to ask the questions you want

Why did arthel Neville leave Fox News? - Answers

Jul 7, 2025 · Copyright ©2025 Answers.com. All Rights Reserved. The material on this site can not be reproduced, distributed, transmitted, cached or otherwise used, except with prior written ...

What is number 1000000000000000000000000000000 in words

Mar 31, 2025 · Oh, that's a big number! Let's paint a picture with words: one followed by 30 zeros is called "one nonillion." Isn't that a lovely word to describe such a vast number? Just imagine ...

Is scottie Scott of the whispers married? - Answers

Aug 29, 2023 · JWalter Scott's wife, Charlotte Margaret Carpenter, was of English descent. She was born in 1771 and married Scott in 1797. Their family background and social status were ...

Why did Brian Alvey divorce? - Answers

Mar 27, 2025 · Brian Alvey divorced due to personal differences and challenges in their relationship, as is common in many marriages. While specific details about the reasons for ...

IS 700 fema course - Answers

May 28, 2025 · Those who have taken the FEMA IS-235 course will have to obtain the answers for the Emergency Planning test through studying the information provided during the course. ...

Who are the female cast members of tmz? - Answers

Feb 11, 2025 · Oh honey, let me break it down for you. The female cast members of TMZ include the fabulous Raquel Harper, the fierce Van Lathan, and the sassy Anna Kachikyan. These ...

Does Oscar blaketon die on heartbeat? - Answers

Jan 5, 2023 · Copyright ©2025 Answers.com. All Rights Reserved. The material on this site can not be reproduced, distributed, transmitted, cached or otherwise used, except with prior written ...

What state in Mexico contains Mexico City? - Answers

Sep 1, 2023 · Copyright ©2025 Answers.com. All Rights Reserved. The material on this site can not be reproduced, distributed, transmitted, cached or otherwise used, except with prior written ...

What are Kasey Annabelle and October Gymnasts models full ...

Mar 26, 2025 · Kasey Annabelle's full name is Kasey Annabelle Lutz, while October Gymnast's full name is October Grace. Both are athletes known for their accomplishments in gymnastics. ...

Answers - The Most Trusted Place for Answering Life's Questions

Answers is the place to go to get the answers you need and to ask the questions you want

Why did arthel Neville leave Fox News? - Answers

Jul 7, 2025 · Copyright ©2025 Answers.com. All Rights Reserved. The material on this site can not be reproduced, distributed, transmitted, cached or otherwise used, except with prior written ...

What is number 1000000000000000000000000000000 in words

Mar 31, 2025 · Oh, that's a big number! Let's paint a picture with words: one followed by 30 zeros is called "one nonillion." Isn't that a lovely word to describe such a vast number? Just imagine ...

Is scottie Scott of the whispers married? - Answers

Aug 29, 2023 · JWalter Scott's wife, Charlotte Margaret Carpenter, was of English descent. She was born in 1771 and married Scott in 1797. Their family background and social status were ...

Why did Brian Alvey divorce? - Answers

Mar 27, 2025 · Brian Alvey divorced due to personal differences and challenges in their relationship, as is common in many marriages. While specific details about the reasons for ...

IS 700 fema course - Answers

May 28, 2025 · Those who have taken the FEMA IS-235 course will have to obtain the answers for the Emergency Planning test through studying the information provided during the course. ...

Who are the female cast members of tmz? - Answers

Feb 11, 2025 · Oh honey, let me break it down for you. The female cast members of TMZ include the fabulous Raquel Harper, the fierce Van Lathan, and the sassy Anna Kachikyan. These ...

Does Oscar blaketon die on heartbeat? - Answers

Jan 5, 2023 · Copyright ©2025 Answers.com. All Rights Reserved. The material on this site can not be reproduced, distributed, transmitted, cached or otherwise used, except with prior written ...

What state in Mexico contains Mexico City? - Answers

Sep 1, 2023 · Copyright ©2025 Answers.com. All Rights Reserved. The material on this site can not be reproduced, distributed, transmitted, cached or otherwise used, except with prior written ...

What are Kasey Annabelle and October Gymnasts models full ...

Mar 26, 2025 · Kasey Annabelle's full name is Kasey Annabelle Lutz, while October Gymnast's full name is October Grace. Both are athletes known for their accomplishments in gymnastics. ...

Unlock the secrets of calculus with our comprehensive guide! Find clear answers to calculus of a single variable. Learn more and enhance your skills today!

[Back to Home](#)