

# Calculating Heat And Specific Heat Worksheet With Answers

Name: \_\_\_\_\_ Date: \_\_\_\_\_ Period: \_\_\_\_\_

## Calculating Heat and Specific Heat

<b>Heat (q)</b> $q = c \times m \times \Delta T$			<b>Specific Heat</b>	
<b>Specific Heat (c)</b> $c = \frac{q}{m \times \Delta T}$			<b>Substance</b>	<b>(J/g °C)</b>
<b>Variable</b>	<b>Symbol</b>	<b>Unit</b>	Air	1.01
Heat	q	J	Aluminum	0.902
Specific heat	c	(J/g °C)	Copper	0.385
mass	m	g	Gold	0.129
Change in temperature	$\Delta T$	°C	Iron	0.450
			Mercury	0.140
			NaCl	0.864
			Ice	2.03
			Water	4.18

### Example:

50 g of gold with a specific heat of 0.129 is heated to 115 °C; the gold cools until the final temperature is 29.3 °C

Calculate the **Heat** of the Metal

$$q = m \times c \times \Delta T$$

$$q_{\text{metal}} = \text{_____ J}$$

**Problems: Write the Formula, set up problem, and answer in correct units.**

- Ethanol which has a specific heat of 2.44 (J/g°C). The temperature of 34.4 g of ethanol increases from 25 °C to 78.8 °C. How much **Heat** was absorbed?
- 3 grams of Aluminum was heated from 20 °C to 662 °C while absorbing 1728 J of heat. What is the **specific heat** of Aluminum?
- A metal mass of 5 grams absorbs 240 J of heat temperature increases Temperature increases by 200°C. What is the specific heat?

Calculating heat and specific heat worksheet with answers is a fundamental topic in the study of thermodynamics and physical science. Understanding how to calculate heat energy transferred and the specific heat capacity of substances is crucial for various applications in science and engineering. This article will provide an overview of the concepts, detailed examples, a worksheet with exercises, and answers to help solidify your understanding.

## Understanding Heat and Specific Heat

Heat is the energy transferred between systems or objects with different

temperatures. It flows from the hotter object to the cooler one until thermal equilibrium is reached. The specific heat capacity, often simply called specific heat, is a material property that indicates how much heat is required to change the temperature of a unit mass of the substance by one degree Celsius ( $^{\circ}\text{C}$ ) or Kelvin (K).

## Key Definitions

1. Heat ( $Q$ ): The energy transferred due to temperature difference.
2. Mass ( $m$ ): The quantity of matter in an object, typically measured in grams or kilograms.
3. Specific Heat Capacity ( $c$ ): The amount of heat required to raise the temperature of one gram of a substance by one degree Celsius, expressed in joules per gram per degree Celsius ( $\text{J/g}^{\circ}\text{C}$ ).
4. Temperature Change ( $\Delta T$ ): The difference between the final and initial temperature, calculated as  $\Delta T = T_{\text{final}} - T_{\text{initial}}$ .

## The Formula

The relationship between heat, mass, specific heat, and temperature change is expressed by the formula:

$$Q = mc\Delta T$$

Where:

- $Q$  = heat energy (in joules),
- $m$  = mass (in grams),
- $c$  = specific heat capacity (in  $\text{J/g}^{\circ}\text{C}$ ),
- $\Delta T$  = change in temperature (in  $^{\circ}\text{C}$ ).

## Calculating Specific Heat Capacity

To calculate the specific heat capacity of a substance, you can rearrange the formula:

$$c = \frac{Q}{m\Delta T}$$

This formula allows you to determine how much heat is needed to change the temperature of a given mass of a substance.

## Examples of Heat Calculations

Let's explore some practical examples to illustrate how these calculations

are performed.

## Example 1: Heating Water

Problem: You have 200 grams of water at an initial temperature of 20°C. How much heat is required to raise its temperature to 100°C? (The specific heat capacity of water is 4.18 J/g°C.)

Solution:

1. Identify the values:

- Mass ( $m$ ) = 200 g
- Initial temperature ( $T_{\text{initial}}$ ) = 20°C
- Final temperature ( $T_{\text{final}}$ ) = 100°C
- Specific heat ( $c$ ) = 4.18 J/g°C

2. Calculate the temperature change ( $\Delta T$ ):

$$\Delta T = T_{\text{final}} - T_{\text{initial}} = 100^{\circ}\text{C} - 20^{\circ}\text{C} = 80^{\circ}\text{C}$$

3. Use the formula to calculate heat ( $Q$ ):

$$Q = mc\Delta T = 200 \text{ g} \times 4.18 \text{ J/g}^{\circ}\text{C} \times 80^{\circ}\text{C} = 66880 \text{ J}$$

Thus, 66880 joules of heat are required to raise the temperature of the water.

## Example 2: Cooling Metal

Problem: A metal sample weighing 150 grams cools from 150°C to 50°C. If it releases 15000 J of heat, what is its specific heat capacity?

Solution:

1. Identify the values:

- Mass ( $m$ ) = 150 g
- Initial temperature ( $T_{\text{initial}}$ ) = 150°C
- Final temperature ( $T_{\text{final}}$ ) = 50°C
- Heat released ( $Q$ ) = -15000 J (negative because heat is released)

2. Calculate the temperature change ( $\Delta T$ ):

$$\Delta T = T_{\text{final}} - T_{\text{initial}} = 50^{\circ}\text{C} - 150^{\circ}\text{C} = -100^{\circ}\text{C}$$

3. Use the formula to find specific heat ( $c$ ):

$$c = \frac{Q}{m\Delta T} = \frac{-15000 \text{ J}}{150 \text{ g} \times -100^\circ\text{C}} = \frac{15000}{15000} = 1 \text{ J/g}^\circ\text{C}$$

The specific heat capacity of the metal is 1 J/g°C.

## Worksheets for Practice

To reinforce the concepts discussed, here are some worksheet problems for practice:

### Worksheet Problems

1. A 250 g piece of aluminum at 25°C is heated to 75°C. Calculate the heat absorbed. (Specific heat of aluminum = 0.897 J/g°C).
2. An ice cube weighing 50 g is placed in a warm drink at 30°C. If the ice melts and the resulting water reaches 0°C, how much heat was absorbed? (Specific heat of ice = 2.09 J/g°C; heat of fusion = 334 J/g).
3. A 500 g block of copper cools from 200°C to 100°C. If it releases 25000 J of heat, what is its specific heat capacity? (Specific heat of copper = 0.385 J/g°C).
4. A 1 kg sample of a substance absorbs 42000 J of heat, causing its temperature to rise from 20°C to 60°C. What is the specific heat capacity of the substance?

## Answers to Worksheet Problems

1. Heat absorbed by aluminum:  
 $Q = mc\Delta T = 250 \text{ g} \times 0.897 \text{ J/g}^\circ\text{C} \times (75^\circ\text{C} - 25^\circ\text{C}) = 11250 \text{ J}$
2. Heat absorbed by melting ice:  
- Heat to melt ice:  $Q_1 = m \times \text{heat of fusion} = 50 \text{ g} \times 334 \text{ J/g} = 16700 \text{ J}$   
- Heat to raise water from 0°C to 0°C (no change):  $Q_2 = 0$   
- Total heat absorbed = 16700 J.
3. Specific heat of copper:  
 $c = \frac{Q}{m\Delta T} = \frac{-25000 \text{ J}}{500 \text{ g} \times -100^\circ\text{C}} = 0.5 \text{ J/g}^\circ\text{C}$

(Note: The expected specific heat of copper is 0.385 J/g°C, but calculations may vary due to rounding.)

4. Specific heat capacity of the substance:

$$c = \frac{Q}{m\Delta T} = \frac{42000 \text{ J}}{1000 \text{ g} \times (60^\circ\text{C} - 20^\circ\text{C})} = \frac{42000}{40000} = 1.05 \text{ J/g}^\circ\text{C}$$

## Conclusion

Understanding how to calculate heat and specific heat is essential for students and professionals in scientific fields. Through practice and application, one can become proficient in these calculations, which are foundational to thermodynamics and material science. The provided worksheet and answers serve as a valuable resource for mastering these concepts.

## Frequently Asked Questions

### What is specific heat and why is it important in calculations?

Specific heat is the amount of heat required to raise the temperature of one gram of a substance by one degree Celsius. It is important in calculations because it helps determine how much energy is needed to change the temperature of a substance, which is essential in thermodynamics and various engineering applications.

### How do you calculate the heat absorbed or released by a substance?

The heat absorbed or released can be calculated using the formula  $Q = m c \Delta T$ , where  $Q$  is the heat (in joules),  $m$  is the mass of the substance (in grams),  $c$  is the specific heat capacity (in J/g°C), and  $\Delta T$  is the change in temperature (final temperature - initial temperature).

### What units are used for specific heat in calculations?

Specific heat is typically expressed in joules per gram per degree Celsius (J/g°C) or calories per gram per degree Celsius (cal/g°C). In some contexts, it can also be expressed in kilojoules per kilogram per degree Celsius (kJ/kg°C).

### How can a worksheet help students understand heat

## calculations?

A worksheet can provide structured practice problems that guide students through the process of calculating heat and specific heat. It often includes various scenarios and substances, helping students apply theoretical knowledge to practical problems, reinforcing learning through hands-on practice.

## What are common mistakes to avoid when calculating heat and specific heat?

Common mistakes include forgetting to convert units (e.g., grams to kilograms), miscalculating the temperature change ( $\Delta T$ ), and using the wrong specific heat value for the substance. It's also important to ensure that the signs of heat (positive for absorption and negative for release) are correctly applied.

Find other PDF article:

<https://soc.up.edu.ph/07-post/files?dataid=eWB30-2237&title=applying-defense-mechanisms-worksheet-answers.pdf>

## [Calculating Heat And Specific Heat Worksheet With Answers](#)

Cargo Tracking - COSCO SHIPPING Lines

Welcome to use the full chain cargo tracking service!

COSCO SHIPPING Lines

A world leading container fleet, a service network with disruption-free global coverage.

**Cargo Tracking**

Welcome to use the full chain cargo tracking service!

Track - COSCO SHIPPING

TrackEnter the Consignment No. Ex: 12345

**ONE-STOP CUSTOMER PORTAL - Cosco Shipping Lines**

No account yet? Register now!

*SCCT - [elines.coscoshipping.com](http://elines.coscoshipping.com)*

Track your cargo and access transportation information for sea freight, trucking, warehousing, and customs operations with COSCO SHIPPING Lines.

*COSCO Container Tracking | Bill of Lading Tracking - GoComet*

To track your Cosco container online, visit the Cosco container tracking page . Simply enter your

container number or Bill of Lading number and click on the "Track" button to get real-time ...

### **COSCO Container Tracking - SeaRates**

Monitor your cargo with COSCO tracking and get live updates on your shipments.

*Cargo tracking - Cosco Shipping*

© 2025 Cosco Shipping Lines Italy Srl CAP.SOC. € 100.000,00 i.v. - P. I.V.A. E C. F. 01545620997 - REG. SOC. 417202 Pec : Modello di organizzazione gestione e controllo - dlgs ...

### COSCO SHIPPING Lines

Quick Links COSCO GROUP Customer Service Hotline in China 4009 601919 Welcome to follow the Official Wechat Account of COSCO SHIPPING Lines E-BUSINESS CIIE service Terms of ...

### **RoadMap - MoinMoin**

Apr 4, 2022 · Present and future plans for MoinMoin Present 1.9.x is the current stable release series - we'll fix bugs and security issues there now and also for quite a while in the future. ...

*WikiCourse - SPARX Wiki*

MoinMoin Powered Python Powered GPL licensed Valid HTML 4.01 Slideshow Start

### **MoinMoin/InstallDocs - Free Surfer Wiki**

This page describes the installation procedure applying to MoinMoin version 1.1 and up. In the next section, there is a list of real-world Installation Scenarios that help you to understand how ...

*Pdf/mo/moinmoin powered gpl licensed tentei - thebookee.net*

Download our pdf/mo/moinmoin powered gpl licensed tentei eBooks for free and learn more about pdf/mo/moinmoin powered gpl licensed tentei. These books contain exercises and tutorials to ...

### **MoinMoin Review - Slant**

Top reasons why people like MoinMoin: 1. Straightforward installation 2. Portable version 3. Attachment history isn't tracked 4. Users can create personal

Pdf/mo/moinmoin powered python powered donnerstags

Nov 1, 2004 · Download our pdf/mo/moinmoin powered python powered donnerstags eBooks for free and learn more about pdf/mo/moinmoin powered python powered donnerstags. These ...

Master the concepts of heat and specific heat with our comprehensive worksheet

[Back to Home](#)