

# Calculus Related Rates Formulas

## Step 2. Find applicable formulas

Formulas

$$C = 2\pi r$$

$$A = \pi r^2$$

## Step 3. Identify "Given," "When" and "Need"

"Given"

$$C' = 20$$

"When"

$$C = 10\pi$$

"Need"

$$A' ?$$

## Step 4. Use formulas for all known and unknown variables

$$C' = 2\pi r'$$

$$C = 2\pi r$$

$$A' = 2\pi(r)(r')$$

## Step 5. Plug in the values for the variables

$$20 = 2\pi r'$$

$$10\pi = 2\pi r$$

$$A' = \pi(5)\left(\frac{10}{\pi}\right)$$

## Step 6. Simplify

$$r' = \frac{10}{\pi}$$

$$r = 5$$

$$A' = 50$$

## Step 7. Don't forget to label!

$$A' = 50 \text{ ft}^2/\text{minute}$$

**Calculus related rates formulas** are a crucial part of differential calculus, particularly in real-world applications where quantities change over time. These formulas enable us to determine the rate at which one quantity changes in relation to another. In this article, we will explore the fundamental concepts behind related rates, provide several key examples, and outline the step-by-step process for solving related rates problems.

## Understanding Related Rates

Related rates problems typically involve two or more variables that are changing with respect to time. The relationship between these variables is often defined by a mathematical equation. When one variable changes, it can affect the rate of change of another variable. Calculus related rates formulas help us quantify these changes.

## Key Concepts

1. **Variables and Parameters:** In related rates problems, we often deal with variables that represent quantities, such as height, radius, or volume. Parameters may refer to constants involved in the relationships.
2. **Derivatives:** The core of related rates lies in the use of derivatives, specifically the concept of differentiation with respect to time. When we differentiate variables that depend on time, we can express their rates of change.
3. **Chain Rule:** The chain rule is essential in related rates, allowing us to relate the change in one variable to the change in another through their functional relationships.

## Steps to Solve Related Rates Problems

To effectively solve related rates problems, follow these systematic steps:

1. **Identify the variables:** Determine which quantities are changing and what relationships exist between them.
2. **Write down the equation:** Establish the equation that relates the different variables. This often involves geometry or physics principles.
3. **Differentiate with respect to time:** Apply implicit differentiation to the equation you set up in the previous step, ensuring to use the chain rule appropriately.
4. **Substitute known values:** Insert the values that you know into the differentiated equation, including rates of change and other fixed quantities.
5. **Solve for the unknown rate:** Isolate the variable representing the rate you need to find and solve the equation.

## Examples of Related Rates Problems

To illustrate how to apply calculus related rates formulas, let's examine several examples.

## Example 1: Volume of a Balloon

Problem Statement: A spherical balloon is being inflated. If the radius of the balloon is increasing at a rate of 2 cm/min, at what rate is the volume of the balloon increasing when the radius is 5 cm?

Solution Steps:

1. Identify the variables:

- Let  $(r)$  be the radius (in cm) and  $(V)$  be the volume (in  $\text{cm}^3$ ).

2. Write down the equation:

- The volume of a sphere is given by  $(V = \frac{4}{3} \pi r^3)$ .

3. Differentiate with respect to time:

- Using the chain rule:

$$\frac{dV}{dt} = \frac{dV}{dr} \cdot \frac{dr}{dt} = 4 \pi r^2 \cdot \frac{dr}{dt}$$

4. Substitute known values:

- Given  $(\frac{dr}{dt} = 2)$  cm/min and  $(r = 5)$  cm:

$$\frac{dV}{dt} = 4 \pi (5^2) \cdot 2 = 4 \pi (25) \cdot 2 = 200 \pi \text{ cm}^3/\text{min}$$

5. Rate of volume increase:

- The volume is increasing at a rate of  $(200 \pi \text{ cm}^3/\text{min})$  when the radius is 5 cm.

## Example 2: A Ladder Against a Wall

Problem Statement: A 10-foot ladder is leaning against a wall. If the bottom of the ladder is sliding away from the wall at a rate of 1 ft/sec, how fast is the top of the ladder descending when the bottom is 6 feet from the wall?

Solution Steps:

1. Identify the variables:

- Let  $(x)$  be the distance from the wall to the bottom of the ladder,  $(y)$  be the height of the ladder on the wall, and  $(L = 10)$  ft be the length of the ladder.

2. Write down the equation:

- Using the Pythagorean theorem, we have:

$$x^2 + y^2 = L^2$$

$$x^2 + y^2 = L^2$$

3. Differentiate with respect to time:

- Differentiating both sides gives:

$$2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$$

4. Substitute known values:

- When  $(x = 6)$  ft,  $(L = 10)$  ft, we can find  $(y)$ :

$$6^2 + y^2 = 10^2 \implies 36 + y^2 = 100 \implies y^2 = 64 \implies y = 8$$

ft

- Now substitute  $(\frac{dx}{dt} = 1)$  ft/sec:

$$2(6)(1) + 2(8)\frac{dy}{dt} = 0$$

$$12 + 16\frac{dy}{dt} = 0 \implies 16\frac{dy}{dt} = -12 \implies \frac{dy}{dt} = -\frac{12}{16} = -\frac{3}{4} \text{ ft/sec}$$

5. Rate of descent:

- The top of the ladder is descending at a rate of  $(\frac{3}{4})$  ft/sec when the bottom is 6 feet from the wall.

## Common Formulas Used in Related Rates Problems

Here are some common formulas that often appear in related rates scenarios:

- **Volume of a Sphere:**  $(V = \frac{4}{3} \pi r^3)$
- **Surface Area of a Sphere:**  $(S = 4 \pi r^2)$
- **Volume of a Cylinder:**  $(V = \pi r^2 h)$
- **Area of a Circle:**  $(A = \pi r^2)$
- **Pythagorean Theorem:**  $(a^2 + b^2 = c^2)$

# Conclusion

Calculus related rates formulas serve as a powerful tool for solving problems where two or more quantities change over time. By following a structured approach, utilizing differentiation, and applying the chain rule, we can successfully analyze and calculate rates of change in various contexts. Whether you're dealing with the geometry of shapes or the dynamics of moving objects, related rates provide essential insights that are applicable in fields ranging from physics to engineering. Understanding these concepts not only enhances your calculus skills but also prepares you for real-world applications of mathematics.

## Frequently Asked Questions

### What are related rates in calculus?

Related rates are a type of problem in calculus that involve finding the rate at which one quantity changes in relation to another. These problems typically involve differentiating equations that relate two or more variables.

### How do you set up a related rates problem?

To set up a related rates problem, start by identifying the quantities that are changing and how they are related. Then, write an equation that relates these quantities, differentiate both sides with respect to time, and substitute known values to solve for the desired rate.

### What is the formula used for related rates involving volume of a sphere?

The volume  $V$  of a sphere is given by the formula  $V = \frac{4}{3}\pi r^3$ . To find the rate of change of volume with respect to time, differentiate this equation with respect to time:  $dV/dt = 4\pi r^2(dr/dt)$ , where  $dr/dt$  is the rate of change of the radius.

### Can you give an example of a related rates problem?

Sure! An example is finding the rate at which the water level in a conical tank is rising as water is poured into it. If the radius and height of the cone are related, you can use the volume formula  $V = \frac{1}{3}\pi r^2h$  and find  $dr/dt$  in terms of  $dh/dt$  using related rates.

### What is the importance of the Chain Rule in related rates problems?

The Chain Rule is crucial in related rates problems because it allows you to

differentiate composite functions. When quantities are changing with respect to time, the Chain Rule helps relate the rates of change of different variables involved in the problem.

Find other PDF article:

<https://soc.up.edu.ph/24-mark/Book?ID=QwA72-9298&title=gabriel-woman-robin-schone-read-online.pdf>

## Calculus Related Rates Formulas

Calculus -

limit 1629 ...

*thomas calculus 13. baskı türkçe pdf olarak - DonanımHaber Forum*

Sep 7, 2023 · Thomas Calculus'un 13. baskısı, Türkçe PDF formatında çevrimiçi olarak bulunabilir. Bu baskı, öğrencilere kalkülüsün temel kavramlarını anlamada yardımcı olacak ...

calculus vs calculation | WordReference Forums

Aug 10, 2014 · Calculus is a specific and complex branch of mathematics. When used as a metaphor, calculus means the same as calculation but suggests a high degree of complexity ...

calculus [non-mathematical] | WordReference Forums

May 26, 2022 · Calculus is defined as "A particular method or system of calculation or reasoning." I've come across other variants such as national calculus (example: The terrorist attacks ...

Lambda calculus: -

$\lambda$ -calculus, Lambda calculus (Wikipedia) .

Lambda calculus: -

~ , . : $\lambda$  ...

lambda calculus? -

C-H term rewriting system, equational reasoning ...

*THOMAS CALCULUS 1-2 TÜRKÇE PDF | DonanımHaber Forum » ...*

Üniversite öğrencileri için Thomas Calculus 1-2 ders kitaplarının Türkçe PDF sürümlerini indirin. Bu kitaplar, kalkülüs kavramlarını Türkçe öğrenmenize yardımcı olacak şekilde özel olarak ...

Calculus (dental)

Mar 2, 2021 · 8515 200,000,000 ...

