

# Occupational Therapy Dementia Case Study

ARTICLE

## How can occupational therapy improve the daily performance and communication of an older patient with dementia and his primary caregiver?

*A case study*

MAUD J.L. GRAFF *Radboud University Nijmegen  
Medical Centre, The Netherlands*

MYRRA J.M. VERNOOIJ-DASSEN *Radboud University Nijmegen  
Medical Centre, The Netherlands*

JANA ZAJEC *Radboud University Nijmegen  
Medical Centre, The Netherlands*

MARCEL G.M. OLDE-RIKKERT *Radboud University Nijmegen  
Medical Centre, The Netherlands*

WILLIBRORD H.L. HOEFNAGELS *Radboud University Nijmegen  
Medical Centre, The Netherlands*

JOOST DEKKER *VU University Amsterdam,  
The Netherlands*



### Abstract

**Objective:** To enhance insight into the process of occupational therapy (OT) and the changes after OT, in an older patient with mild dementia and his primary caregiver.

**Design and setting:** Case study: content analysis of an OT patient record.

**Intervention:** System-based OT at home using a guideline focusing on both patient's performance in daily activities and caregiver's cognition on patient behaviour and caregiver role and focusing on adaptation of the physical environment.

DOI: 10.1177/1471301206069918

Downloaded from dem.sagepub.com at Radboud Universiteit Nijmegen on January 5, 2015

Occupational therapy dementia case study is a critical area of study that highlights the importance of therapeutic interventions in enhancing the quality of life for individuals diagnosed with dementia. As dementia progresses, it impairs cognitive functions, leading to challenges in daily living activities. Occupational therapy (OT) plays a vital role in providing strategies and techniques to help individuals maintain independence and engage in meaningful activities. This article will explore a detailed case study of a patient with dementia, the therapeutic interventions used, and the outcomes observed.

# Understanding Dementia and Its Impact

Dementia encompasses a range of symptoms that affect memory, thinking, and social abilities. It is not a specific disease but rather a general term that describes a decline in cognitive function severe enough to interfere with daily life. The most common form of dementia is Alzheimer's disease, but there are several other types, including vascular dementia, Lewy body dementia, and frontotemporal dementia.

## Common Symptoms of Dementia

- Memory loss, particularly short-term memory
- Difficulty in performing familiar tasks
- Challenges in communication and language
- Disorientation to time and place
- Changes in mood and personality
- Poor judgment and decision-making

As these symptoms progress, individuals may require assistance with daily living activities, which is where occupational therapy can make a significant difference.

## Case Study Overview

This case study focuses on Mrs. Jane Smith, a 75-year-old woman diagnosed with moderate Alzheimer's disease. She lived alone in her home and had been experiencing increasing difficulties with daily tasks, including cooking, managing medications, and personal hygiene.

## Initial Assessment

Upon referral to occupational therapy, an initial assessment was conducted to determine Mrs. Smith's abilities, challenges, and goals. The assessment included:

1. Cognitive Evaluation: This involved standardized tests to assess memory, attention, and problem-solving skills.
2. Functional Assessment: Evaluating her ability to perform activities of daily living (ADLs), such as dressing, bathing, and meal preparation.
3. Environmental Assessment: Observing her living environment for safety hazards and accessibility issues.
4. Interview: Engaging in a conversation with Mrs. Smith and her family to understand her interests, preferences, and social support systems.

## Identified Challenges

The assessment revealed several challenges that Mrs. Smith faced:

- Memory Impairment: Difficulty remembering appointments and taking medications.
- Safety Concerns: Increased risk of falls due to clutter and poor lighting in her home.
- Difficulty with ADLs: Struggled with bathing and dressing independently.
- Social Isolation: Reduced social interactions and engagement in community activities.

## Goals of Occupational Therapy Intervention

Based on the assessment, specific goals were established for Mrs. Smith's occupational therapy intervention:

1. Enhance Independence: Improve her ability to perform ADLs independently.
2. Increase Safety: Modify her environment to reduce safety hazards.
3. Support Cognitive Function: Implement strategies to improve memory and cognitive functioning.
4. Promote Social Engagement: Encourage participation in social activities and community programs.

## Therapeutic Interventions

The occupational therapy intervention plan included a combination of strategies tailored to meet Mrs. Smith's needs.

### 1. Environmental Modifications

- Decluttering: Organized her living space to minimize hazards, allowing for easier navigation.
- Lighting Adjustments: Improved lighting in key areas, such as hallways and the kitchen, to enhance visibility.
- Labeling: Used labels on essential items (e.g., refrigerator, bathroom) to aid memory and promote independence.

### 2. Cognitive Strategies

- Memory Aids: Introduced the use of a daily planner and medication reminder system to help manage appointments and medications.

- Routine Establishment: Worked with Mrs. Smith to establish a consistent daily routine that included set times for meals, activities, and self-care.
- Engagement in Cognitive Activities: Encouraged participation in puzzles, reading, and other brain-stimulating activities to support cognitive function.

### **3. Skills Training for ADLs**

- Personal Care Techniques: Provided hands-on training for bathing and dressing, breaking down tasks into manageable steps.
- Meal Preparation: Developed simple, nutritious meal plans and cooking techniques that Mrs. Smith could easily follow.
- Adaptive Equipment: Recommended tools such as a reacher, non-slip mats, and adaptive utensils to promote safety and independence during ADLs.

### **4. Social Engagement Activities**

- Community Programs: Assisted Mrs. Smith in finding local community centers offering social activities for seniors.
- Family Involvement: Educated her family about the importance of social engagement and encouraged regular visits and outings.
- Group Therapy Sessions: Suggested participation in group therapy sessions focused on reminiscence therapy to stimulate conversation and social interaction.

## **Outcomes and Progress Monitoring**

The progress of Mrs. Smith was monitored regularly through follow-up assessments and family feedback.

### **Progress Indicators**

- Improved Independence: After six weeks of intervention, Mrs. Smith reported being able to dress and bathe herself with minimal assistance.
- Increased Safety: Family members noted a significant reduction in safety concerns due to environmental modifications.
- Enhanced Cognitive Function: Mrs. Smith demonstrated improved memory recall when using the daily planner, and she successfully managed her medications.
- Social Engagement: She began attending community events and participating in group activities,

significantly reducing her feelings of isolation.

## Conclusion

The occupational therapy dementia case study of Mrs. Jane Smith illustrates the profound impact that targeted therapeutic interventions can have on the lives of individuals with dementia. Through a comprehensive assessment, personalized intervention strategies, and ongoing support, Mrs. Smith was able to regain a sense of independence and improve her overall quality of life.

Occupational therapists play a crucial role in helping individuals with dementia navigate the challenges of daily living, emphasizing the importance of maintaining autonomy and engaging in meaningful activities. As the population ages, the demand for occupational therapy services for dementia will continue to grow, underscoring the need for ongoing research and development of effective therapeutic strategies in this field.

By focusing on the individual's strengths and providing tailored interventions, occupational therapy can empower those living with dementia to lead fulfilling lives, despite the challenges they face.

## Frequently Asked Questions

### **What are the primary goals of occupational therapy for dementia patients?**

The primary goals include enhancing the patient's independence in daily activities, improving quality of life, maintaining cognitive function, and providing strategies for managing behavioral symptoms.

### **How can occupational therapists assess the needs of a dementia patient?**

Occupational therapists can assess the needs through standardized assessments, interviews with the patient and caregivers, observational assessments, and evaluating the patient's home environment.

### **What types of interventions might be used in occupational therapy for dementia?**

Interventions may include cognitive rehabilitation, sensory stimulation, adaptive equipment training, environmental modifications, and caregiver education.

## **How does occupational therapy support caregivers of dementia patients?**

Occupational therapy provides caregivers with training, resources, and emotional support, helping them to better manage care demands and reduce stress.

## **What role does activity engagement play in occupational therapy for dementia?**

Activity engagement helps maintain cognitive and physical abilities, promotes social interaction, and can reduce behavioral symptoms through meaningful and familiar activities.

## **How often should occupational therapy sessions be scheduled for dementia patients?**

The frequency of sessions can vary based on the patient's needs, but typically ranges from once a week to multiple times a week, as determined by the therapist.

## **What is the impact of customizing therapy plans for individual dementia patients?**

Customizing therapy plans ensures that interventions are relevant to the patient's interests, abilities, and stage of dementia, leading to more effective outcomes and greater patient engagement.

## **What are common barriers to effective occupational therapy for dementia patients?**

Common barriers include communication difficulties, resistance to change, physical limitations, and environmental challenges, which therapists must navigate to provide effective care.

## **How can technology be incorporated into occupational therapy for dementia?**

Technology can be used through memory aids, interactive applications for cognitive training, virtual reality for engagement, and telehealth services to reach patients in remote areas.

Find other PDF article:

<https://soc.up.edu.ph/49-flash/files?ID=gJS55-4672&title=python-coding-interview-cheat-sheet.pdf>

## **[Occupational Therapy Dementia Case Study](#)**

### Occupational health

Jun 12, 2025 · Occupational health is an area of work in public health to promote and maintain highest degree of physical, mental and social well-being of workers in all occupations. Its ...

### **Occupational hazards in the health sector : Training slides**

Mar 6, 2025 · Overview Following the publication of our Caring for those who care: Guide for the development and implementation of occupational health and safety programmes for health ...

### **Occupational hazards in the health sector**

This e-tool is intended for use by people in charge of occupational health and safety for health workers at the national, subnational and facility levels.

### **A life in motion: Lyle's journey as an Occupational Therapist**

Mar 26, 2025 · Occupational therapy is a client-centred health profession concerned with promoting health and wellbeing through occupation. The primary goal of occupational therapy ...

### *Occupational health: health workers - World Health Organization ...*

Nov 7, 2022 · Safeguarding the health, safety and well-being of health workers is fundamental for well-functioning and resilient health systems. Health workers face a range of occupational risks ...

### Occupational injuries - World Health Organization (WHO)

Occupational injuries Common injuries among health workers are slips, trips and falls, road traffic injuries (ambulance crashes, motorbike and bicycle injuries), electric shock, explosions and fire.

### *Burn-out an "occupational phenomenon": International ...*

May 28, 2019 · Burn-out is included in the 11th Revision of the International Classification of Diseases (ICD-11) as an occupational phenomenon. It is not classified as a medical condition. It ...

### Guidelines for HIV post-exposure prophylaxis

Jul 22, 2024 · WHO's updated PEP guidelines prioritize broader access to PEP, including community-based delivery and task sharing to mitigate barriers such as stigma and to ensure ...

### *Occupational health*

Jun 1, 2023 · Occupational health deals with all aspects of health and safety in the workplace and has a strong focus on primary prevention of hazards. The health of the workers has several ...

### Protecting health and safety of health workers

Providing occupational health and safety measures for protecting health workers and is also fundamental for well-functioning and resilient health systems, quality of care and maintaining a ...

### *Occupational health*

Jun 12, 2025 · Occupational health is an area of work in public health to promote and maintain highest degree of physical, mental and social well-being of workers in all occupations. Its ...

### **Occupational hazards in the health sector : Training slides**

Mar 6, 2025 · Overview Following the publication of our Caring for those who care: Guide for the development and implementation of occupational health and safety programmes for health ...

### **Occupational hazards in the health sector**

This e-tool is intended for use by people in charge of occupational health and safety for health workers at the national, subnational and facility levels.

## **A life in motion: Lyle's journey as an Occupational Therapist**

Mar 26, 2025 · Occupational therapy is a client-centred health profession concerned with promoting health and wellbeing through occupation. The primary goal of occupational therapy ...

## Occupational health: health workers - World Health Organization ...

Nov 7, 2022 · Safeguarding the health, safety and well-being of health workers is fundamental for well-functioning and resilient health systems. Health workers face a range of occupational ...

## **Occupational injuries - World Health Organization (WHO)**

Occupational injuries Common injuries among health workers are slips, trips and falls, road traffic injuries (ambulance crashes, motorbike and bicycle injuries), electric shock, explosions and fire.

## **Burn-out an "occupational phenomenon": International ...**

May 28, 2019 · Burn-out is included in the 11th Revision of the International Classification of Diseases (ICD-11) as an occupational phenomenon. It is not classified as a medical ...

## **Guidelines for HIV post-exposure prophylaxis**

Jul 22, 2024 · WHO's updated PEP guidelines prioritize broader access to PEP, including community-based delivery and task sharing to mitigate barriers such as stigma and to ensure ...

## **Occupational health**

Jun 1, 2023 · Occupational health deals with all aspects of health and safety in the workplace and has a strong focus on primary prevention of hazards. The health of the workers has several ...

## Protecting health and safety of health workers

Providing occupational health and safety measures for protecting health workers and is also fundamental for well-functioning and resilient health systems, quality of care and maintaining a ...

Explore our in-depth occupational therapy dementia case study to understand effective strategies and interventions. Discover how therapy can enhance patient care today!

[Back to Home](#)