




Ozempic Dispensing Quick Reference Guide

Ozempic® Dispensing

Quick Reference Guide

OZEMPIC®

semaglutide injection 0.25mg, 1mg, 2mg

	Ozempic® 0.25-mg and 0.5-mg dose pen pack Delivers 0.25 and 0.5 mg doses only			Ozempic® 1-mg dose pen pack Delivers 1 mg dose only		Ozempic® 2-mg dose pen pack Delivers 2 mg dose only	
Trade Pack							
NDC	0169-4181-13			0169-4130-13		0169-4772-13	
Days Supply	Sample or Initial (42 days)	1 Month (28 days)	3 Months (84 days)	1 Month (28 days)	3 Months (84 days)	1 Month (28 days)	3 Months (84 days)
Intent of Prescription	Sample or initial prescription for new starts	1-month prescription for maintenance on 0.5 mg	3-month prescription for maintenance on 0.5 mg	1-month prescription for maintenance on 1 mg	3-month prescription for maintenance on 1 mg	1-month prescription for maintenance on 2 mg	3-month prescription for maintenance on 2 mg
Strength	2 mg per 3 mL (0.68 mg/mL)			4 mg per 3 mL (1.34 mg/mL)		8 mg per 3 mL (2.68 mg/mL)	
Dosage Form	Solution			Solution		Solution	
SIG	Sample or Initial Rx: 0.25 mg SC once weekly for 4 weeks, then 0.5 mg SC once weekly for 2 weeks	Maintenance Rx: Inject 0.5 mg SC once weekly for 4 weeks	Maintenance Rx: Inject 0.5 mg SC once weekly for 12 weeks	Maintenance Rx: Inject 1 mg SC once weekly for 4 weeks	Maintenance Rx: Inject 1 mg SC once weekly for 12 weeks	Maintenance Rx: Inject 2 mg SC once weekly for 4 weeks	Maintenance Rx: Inject 2 mg SC once weekly for 12 weeks
Dispense Quantity	3 mL			3 mL		3 mL	
Needles	6 included			12 included		12 included	
Number of Boxes	1 box			1 box		1 box	

For adults with T2D, along with diet and exercise, to improve glycemic control. For adults with T2D and established CVD, to reduce risk of MACE.

Ozempic® pen packs are not interchangeable. Always ensure prescriptions and dispensing quantities match the intended dosage.

Ozempic dispensing quick reference guide is an essential resource for healthcare professionals and patients alike. As a medication used for the management of Type 2 diabetes, Ozempic (semaglutide) has gained popularity due to its efficacy in lowering blood sugar levels and promoting weight loss. This guide aims to provide comprehensive information on dispensing Ozempic, including dosage, administration, storage, and patient education.

Understanding Ozempic

Ozempic is a GLP-1 receptor agonist that mimics the incretin hormones, which help regulate blood sugar levels. It is administered as a subcutaneous injection and is typically prescribed to patients who struggle to control their blood sugar levels with diet and exercise alone.

Indications for Use

Ozempic is indicated for:

- Adults with Type 2 diabetes mellitus as an adjunct to diet and exercise to improve glycemic control.
- Weight management in addition to a reduced-calorie diet and increased physical activity for adults who are overweight or obese.

Dosage and Administration

Ozempic is available in pre-filled pens containing 1.5 mL of solution, and it is typically dosed once weekly.

- Initial dose: 0.25 mg once weekly for the first four weeks.
- Maintenance dose: After four weeks, the dose may be increased to 0.5 mg once weekly.
- Further increase: If additional glycemic control is needed, the dose may be increased to 1 mg once weekly after at least four weeks at the 0.5 mg dose.

Dispensing Ozempic

When dispensing Ozempic, it is important to ensure that the patient understands how to use the medication properly. Here are key points to cover:

Storage Requirements

- Ozempic should be stored in the refrigerator between 36°F to 46°F (2°C to 8°C) before first use.
- After the first use, it can be kept at room temperature (up to 86°F or 30°C) for up to 56 days.
- Avoid exposing the pen to extreme temperatures or direct sunlight.

Preparation for Administration

Before administering Ozempic, the following steps should be taken:

1. Gather all necessary supplies, including the Ozempic pen, alcohol swabs, and a sharps container for disposal.
2. Wash hands thoroughly.
3. Inspect the pen for any visible particles or discoloration—if present, do not use the pen.
4. Attach a new needle to the pen.
5. Prime the pen by dialing to 0.25 mg and pressing the injection button until a drop appears at the needle tip.

Injection Technique

Patients should be instructed on proper injection techniques:

- Choose an injection site: abdomen, thigh, or upper arm.
- Clean the injection site with an alcohol swab and let it dry.
- Pinch the skin and insert the needle at a 90-degree angle.
- Press the injection button and hold for a few seconds to ensure complete delivery.
- Dispose of the needle in a sharps container immediately after use.

Patient Education

Effective patient education is vital for the successful use of Ozempic. Healthcare providers should cover the following topics:

Potential Side Effects

Patients should be aware of possible side effects, which may include:

- Nausea
- Vomiting
- Diarrhea
- Constipation
- Abdominal pain
- Risk of pancreatitis

Patients should be advised to report severe abdominal pain or persistent nausea and vomiting to their healthcare provider.

Monitoring and Follow-Up

Regular follow-up appointments are essential to monitor the patient's response to Ozempic. Key aspects to discuss during these visits include:

- Blood sugar level monitoring
- Weight management progress
- Potential adjustments to the medication dosage
- Evaluation for any side effects

Diet and Lifestyle Recommendations

Patients should be encouraged to maintain a healthy diet and engage in regular physical activity. Suggestions include:

- Following a balanced diet rich in fruits, vegetables, whole grains, and lean proteins.
- Limiting intake of sugary foods and beverages.
- Engaging in at least 150 minutes of moderate-intensity exercise each week.

Conclusion

The **Ozempic dispensing quick reference guide** serves as a valuable tool for healthcare professionals and patients in the management of Type 2 diabetes. Proper understanding of the medication, its administration, and ongoing patient education can significantly enhance treatment outcomes. By adhering to the guidelines outlined in this article, healthcare providers can ensure safe and effective use of Ozempic, ultimately improving the quality of life for their patients.

By equipping patients with the knowledge they need to manage their diabetes effectively, we can foster a more informed and proactive approach to their health care journey. Always remember that open communication between patients and healthcare professionals is key to achieving optimal diabetes management.

Frequently Asked Questions

What is Ozempic and what is it used for?

Ozempic is a prescription medication used to improve blood sugar control in adults with type 2 diabetes, along with diet and exercise.

How is Ozempic administered?

Ozempic is administered as a subcutaneous injection, typically once a week, in the thigh, abdomen, or upper arm.

What should be included in the Ozempic dispensing quick reference guide?

The guide should include dosing information, storage instructions, administration techniques, side effects, and patient counseling points.

Are there any specific storage requirements for Ozempic?

Ozempic should be stored in the refrigerator and can be kept at room temperature for up to 56 days after the first use.

What are common side effects of Ozempic?

Common side effects include nausea, vomiting, diarrhea, abdominal pain, and constipation.

Can Ozempic be used in conjunction with other diabetes medications?

Yes, Ozempic can be used with other diabetes medications, but patients should consult their healthcare provider for personalized advice.

What should a patient do if they miss a dose of Ozempic?

If a patient misses a dose, they should take it as soon as they remember, but skip it if it's close to the time of their next dose; never take two doses at once.

Is there a specific patient population that should avoid Ozempic?

Ozempic is not recommended for individuals with a personal or family history of medullary thyroid carcinoma or Multiple Endocrine Neoplasia syndrome type 2.

Find other PDF article:

<https://soc.up.edu.ph/60-flick/Book?docid=axm46-9473&title=the-modern-art-of-chinese-cooking.pdf>

[Ozempic Dispensing Quick Reference Guide](#)

Ozempic | Diabetes UK Forum

Jan 10, 2022 · started Ozempic (semaglutide) .25mg on Friday , Saturday I experienced extreme dizziness and nausea with vomiting, it really floored me for 24hrs. contacted my diabetic clinic ...

Nach über einem Jahr Ozempic - diabetes-forum.de

May 8, 2025 · Hallo zusammen ich bin Bernhard und habe jetzt fast eineinhalb Jahre Ozempic hinter mir. Meine aktuelle Wochendosis beträgt 1mg. Nach anfänglichen Schwierigkeiten wie ...

Ozempic | Diabetes UK Forum

Nov 22, 2020 · Hello, i'm new here but i have been active in diabetes online elsewhere so there's a few familiar faces. I've read a few threads and there seems to be people asking questions on ...

Ozempic | Diabetes UK Forum

Jan 25, 2024 · Hi, just a bit of advice if possible? I have been in the pre diabetic range for as long as I can remember (I'm now 59) out of five siblings four of us have diabetes. HbA1c had ...

Wieviele Dosen sind wirklich im Pen? - diabetes-forum

Mar 2, 2021 · Hallo zusammen, ich nehme seit 4 Wochen Ozempic 0,25 mg wegen einer Insulinresistenz und zur Gewichtsabnahme. Auf der Packung des Pen steht '4 Dosen', ich ...

diabetes-forum.de // Forum // Allgemein // Ozempic

Feb 14, 2024 · Guten Abend, Hier mal meine Erfahrungen mit Ozempic. Die ersten vier Wochen Ozempic 0,25 bis auf Sodbrennen alles gut, wenn ich auf den Rücken geschlafen ...

Erfahrungen mit ozempic - diabetes-forum

Feb 29, 2020 · Guten Tag. Ich nehme ozempic seit ungefähr 2 Monaten. Bin 44 Jahre alt und habe Diabetes Typ 2. Bei der Dose von 0,25 MG war alles gut: In einem Monat gab es ...

ozempic | Diabetes UK Forum

Aug 14, 2019 · I started ozempic a couple of weeks ago, so still on the 0.25 dose. However, the side effects of sulphur belching, flatulence and constant diarrhoea are severe. Do these ...

diabetes-forum.de // Forum // Suchen // Ozempic

Apr 7, 2025 · Hallo Marry, du machst nichts falsch, wenn du das Ozempic weiter gibst. Auch wenn an gemeinhin denkt, dass die Weitergabe von verschreibungspflichtigen Medikamenten ...

Has anybody with Type 1 in the UK been able to take ...

Feb 9, 2024 · On the UK diabetes Type 1 face book group there are a few people that have been prescribed the above, weogvy and ozempic too. That is what prompted me to ask the ...

Ozempic | Diabetes UK Forum

Jan 10, 2022 · started Ozempic (semaglutide) .25mg on Friday, Saturday I experienced extreme dizziness and nausea with vomiting, it really floored me for 24hrs. contacted my diabetic clinic ...

Nach über einem Jahr Ozempic - diabetes-forum.de

May 8, 2025 · Hallo zusammen ich bin Bernhard und habe jetzt fast eineinhalb Jahre Ozempic hinter mir. Meine aktuelle Wochendosis beträgt 1mg. Nach anfänglichen Schwierigkeiten wie ...

Ozempic | Diabetes UK Forum

Nov 22, 2020 · Hello, i'm new here but i have been active in diabetes online elsewhere so there's a few familiar faces. I've read a few threads and there seems to be people asking questions on ...

Ozempic | Diabetes UK Forum

Jan 25, 2024 · Hi, just a bit of advice if possible? I have been in the pre diabetic range for as long as I can remember (I'm now 59) out of five siblings four of us have diabetes. HbA1c had ...

Wieviele Dosen sind wirklich im Pen? - diabetes-forum

Mar 2, 2021 · Hallo zusammen, ich nehme seit 4 Wochen Ozempic 0,25 mg wegen einer Insulinresistenz und zur Gewichtsabnahme. Auf der Packung des Pen steht '4 Dosen', ich ...

diabetes-forum.de // Forum // Allgemein // Ozempic

Feb 14, 2024 · Guten Abend, Hier mal meine Erfahrungen mit Ozempic. Die ersten vier Wochen Ozempic 0,25 bis auf Sodbrennen alles gut, wenn ich auf den Rücken geschlafen habe. Danach ...

Erfahrungen mit ozempic - diabetes-forum

Feb 29, 2020 · Guten Tag. Ich nehme ozempic seit ungefähr 2 Monaten. Bin 44 Jahre alt und habe Diabetes Typ 2. Bei der Dose von 0,25 MG war alles gut: In einem Monat gab es ...

ozempic | Diabetes UK Forum

Aug 14, 2019 · I started ozempic a couple of weeks ago, so still on the 0.25 dose. However, the side effects of sulphur belching, flatulence and constant diarrhoea are severe. Do these ...

[diabetes-forum.de // Forum // Suchen // Ozempic](#)

Apr 7, 2025 · Hallo Marry, du machst nichts falsch, wenn du das Ozempic weiter gibst. Auch wenn an gemeinhin denkt, dass die Weitergabe von verschreibungspflichtigen Medikamenten ...

Has anybody with Type 1 in the UK been able to take ...

Feb 9, 2024 · On the UK diabetes Type 1 face book group there are a few people that have been prescribed the above, weogvy and ozempic too. That is what prompted me to ask the question. ...

"Get your Ozempic dispensing quick reference guide! Discover essential tips and guidelines for safe and effective use. Learn more to optimize your treatment!"

[Back to Home](#)