

Perinatal Mood Disorders Training

REGISTER TODAY
JUNE 18 – 20, 2025

Perinatal Mood & Anxiety Disorders

3-Day Training Presented by Postpartum Support International
Embassy Suites, 300 Court Street, Charleston WV 25301
Registration required

Perinatal Mood Disorders: Components of Care
June 18th-19th
8:00AM – 5:00PM

Participants will be able to identify challenges in diagnosing and treating perinatal mood and anxiety disorders, and describe symptoms of various conditions including depression, anxiety, OCD, PTSD, and psychosis during pregnancy and postpartum.

\$425 Regular Registration
\$325 Student Registration



Advanced Perinatal Mental Health Psychotherapy Training
June 20th
8:00AM – 4:30PM

Participants will be able to understand diagnostic coding related to perinatal mental health (PMH) and discuss evidence-based therapeutic approaches for PMH.

\$250 for Advanced Training
\$150 for Student Registration
You must have completed the two-day course before attending the advanced session

Target Audience: This training is designed for psychiatrists, psychologists, nurses, physicians, social workers, mental health providers, childbirth professionals, social support providers, and anyone in the healthcare community interested in building skills to treat perinatal mood disorders.

Scholarships Available:
Contact Caroline Crislip at ccrislip@wvperinatal.org for more information.

Deadline to register: June 9, 2025



This training is sponsored by the West Virginia Perinatal Partnership and PSI-WV chapter. It is made possible through generous grant support from the WV Department of Human Services.



Postpartum Support International
West Virginia Chapter



West Virginia
Perinatal
PARTNERSHIP

Working together for healthier mothers and babies



Perinatal mood disorders training is a crucial aspect of mental health education for healthcare providers, educators, and support personnel who work with pregnant and postpartum individuals. As awareness of perinatal mood disorders (PMDs) grows, it becomes increasingly important for professionals in various fields to recognize, understand, and address these conditions. This article delves into what perinatal mood disorders are, their prevalence, symptoms, and the importance of specialized training for those working with affected individuals.

Understanding Perinatal Mood Disorders

Definition and Types

Perinatal mood disorders encompass a range of mental health issues that can occur during pregnancy and the postpartum period. The most common types include:

1. **Prenatal Depression:** This occurs during pregnancy and can significantly affect both the mother and the developing fetus.
2. **Postpartum Depression (PPD):** Affects mothers after childbirth, leading to feelings of extreme sadness, anxiety, and exhaustion that can hinder a woman's ability to care for herself and her newborn.
3. **Postpartum Anxiety (PPA):** Involves excessive worry and nervousness after childbirth, which can manifest as panic attacks or obsessive-compulsive behaviors.
4. **Postpartum Psychosis:** A rare but severe condition characterized by hallucinations, delusions, and significant impairment in functioning, requiring immediate medical attention.

Prevalence

Research indicates that about 15-20% of women experience PMDs during the perinatal period. However, this number may be higher due to underreporting. Factors contributing to PMDs include hormonal changes, lack of social support, personal or family history of mental health issues, and stressful life events.

Symptoms of Perinatal Mood Disorders

Recognizing the symptoms of PMDs is essential for early intervention. Common symptoms include:

- Persistent sadness or low mood
- Loss of interest in activities previously enjoyed
- Changes in appetite or weight
- Sleep disturbances (insomnia or excessive sleeping)
- Feelings of guilt or worthlessness
- Difficulty concentrating or making decisions
- Irritability or anger
- Thoughts of self-harm or harming the baby (in severe cases)

It is vital for healthcare providers to be familiar with these symptoms to facilitate timely diagnosis and treatment.

The Importance of Perinatal Mood Disorders Training

1. Enhancing Awareness and Understanding

Training in PMDs helps healthcare professionals:

- Develop a comprehensive understanding of the various types of PMDs and their symptoms.
- Recognize the impact of these disorders on both the mother and child, including potential developmental issues for the child if left untreated.

2. Improving Screening and Diagnosis

Education on PMDs equips providers with the skills to:

- Conduct effective screenings during routine visits, such as the Edinburgh Postnatal Depression Scale (EPDS) or similar tools.
- Differentiate between normal postpartum feelings and clinical depression or anxiety, ensuring appropriate referrals and interventions.

3. Providing Support and Resources

Healthcare providers trained in PMDs can:

- Offer empathy and understanding to affected individuals, reducing stigma and encouraging open discussions about mental health.
- Provide information about local resources, including counseling services, support groups, and educational materials.

4. Promoting Early Intervention and Treatment

Prompt recognition and treatment of PMDs can significantly improve outcomes for mothers and their babies. Training prepares providers to:

- Discuss treatment options, including therapy, medication, and lifestyle changes.
- Develop individualized care plans that involve the mother's healthcare team, ensuring a holistic approach to treatment.

Components of Effective Training Programs

To ensure comprehensive training in perinatal mood disorders, effective programs should include the following components:

1. Evidence-Based Curriculum

Training should be grounded in the latest research, incorporating:

- Current statistics and trends in PMDs.
- Evidence-based practices for screening and treatment.
- Case studies that illustrate the complexities of PMDs.

2. Interactive Learning Experiences

Engaging training methods can enhance retention and application of knowledge. Effective programs may utilize:

- Role-playing scenarios to simulate patient-provider interactions.
- Group discussions to share experiences and strategies for addressing PMDs.
- Workshops on screening tools and treatment modalities.

3. Multidisciplinary Approach

Involving professionals from various fields ensures a comprehensive understanding of PMDs. Training programs can include:

- Psychologists and psychiatrists to discuss mental health treatment.
- Social workers to address community resources and support systems.
- Pediatricians to explore the impact of PMDs on child development.

4. Ongoing Education and Support

Mental health knowledge is continually evolving, so ongoing education is vital. Programs should offer:

- Regular updates on new research and treatment options.

- Opportunities for continued professional development through workshops and conferences.
- Support networks for professionals to share challenges and solutions in addressing PMDs.

Implementing Perinatal Mood Disorders Training

To effectively implement training within healthcare settings, organizations can follow these steps:

1. Assess Training Needs

Conduct surveys or focus groups to identify gaps in knowledge and areas for improvement among staff regarding PMDs.

2. Develop a Training Plan

Create a structured plan that outlines the training objectives, curriculum, and timeline. Ensure that the training is tailored to the specific needs of the organization.

3. Engage Qualified Trainers

Collaborate with mental health professionals who specialize in PMDs to facilitate the training. Their expertise will lend credibility and depth to the program.

4. Evaluate Effectiveness

After training, assess its impact on staff knowledge, confidence, and patient outcomes. Use feedback to refine future training initiatives.

Conclusion

Perinatal mood disorders are significant mental health challenges that affect a large number of individuals during a vulnerable time in their lives. Comprehensive training in PMDs equips healthcare providers with the knowledge and skills necessary to recognize, diagnose, and treat these conditions effectively. By fostering awareness, improving screening practices, and promoting early intervention, we can ensure

better mental health outcomes for mothers and their children. As the landscape of mental health care continues to change, ongoing education and support for professionals in the field remain crucial for combating the stigma surrounding PMDs and providing compassionate care to those in need.

Frequently Asked Questions

What are perinatal mood disorders, and why is training important for healthcare providers?

Perinatal mood disorders include conditions like postpartum depression and anxiety that can affect women during pregnancy and after childbirth. Training is crucial for healthcare providers to recognize symptoms early, provide appropriate support, and improve maternal and infant health outcomes.

What types of training programs are available for perinatal mood disorders?

Training programs for perinatal mood disorders can include online courses, workshops, and certifications focusing on screening, assessment, treatment options, and supportive communication strategies tailored for healthcare professionals.

How can healthcare providers effectively screen for perinatal mood disorders in their patients?

Healthcare providers can use standardized screening tools such as the Edinburgh Postnatal Depression Scale (EPDS) or the Patient Health Questionnaire-9 (PHQ-9) to identify symptoms. Regular follow-ups during prenatal and postnatal visits are also essential for ongoing assessment.

What role does cultural competence play in training for perinatal mood disorders?

Cultural competence is vital in training as it enables healthcare providers to understand and respect diverse backgrounds, beliefs, and experiences, which can influence a patient's experience with perinatal mood disorders and their willingness to seek help.

What are some effective treatment options that health professionals should be aware of when dealing with perinatal mood disorders?

Effective treatment options include psychotherapy, particularly cognitive-behavioral therapy (CBT), pharmacotherapy (antidepressants), and support groups. Training should emphasize a collaborative approach to tailor treatment to individual patient needs.

How can training on perinatal mood disorders impact maternal and infant health outcomes?

Training can significantly improve the ability of healthcare providers to identify and treat perinatal mood disorders, leading to better maternal mental health, enhanced bonding with infants, and reduced risk of developmental issues in children.

Find other PDF article:

<https://soc.up.edu.ph/08-print/pdf?ID=sDU35-8391&title=bachelors-degree-in-civil-engineering.pdf>

Perinatal Mood Disorders Training

Sexual and Reproductive Health and Research (SRH)

Apr 25, 2023 · Maternal and perinatal health Each day, about 800 women across the world die from complications related to pregnancy or childbirth, most of which are preventable or treatable.

Maternal and Infant Health - Public Health Ontario

Jan 20, 2021 · Documents and resources related to the health of mothers and children during pregnancy and the first years of life.

WHO guide for integration of perinatal mental health in maternal ...

Sep 19, 2022 · The guide for integration of perinatal mental health in maternal and child health services outlines an evidence-informed approach describing how program managers, health ...

Evidence Brief: Exploring interventions to address perinatal mental ...

The time from pregnancy to childbirth is commonly referred to as the perinatal period. There are multiple working definitions for 'perinatal period'; however, for the purpose of this Evidence ...

IPC in Perinatology_ENGLISH_2015_Revision - Public Health Ontario

How to cite this document: Ontario Agency for Health Protection and Promotion (Public Health Ontario), Provincial Infectious Diseases Advisory Committee. Best practices for infection ...

Maternal and perinatal death surveillance and response

Maternal and perinatal death surveillance and response (MPDSR) is defined as an essential quality improvement intervention which permits the identification, notification, quantification and ...

Newborn mortality - World Health Organization (WHO)

Mar 14, 2024 · WHO fact sheet on newborn mortality, including key facts, causes, priority strategies, newborn care and WHO response.

Research in maternal, perinatal, and adolescent health

Mar 30, 2019 · WHO develops guidelines on a broad array of clinical, public health, health system, health promotion and implementation strategies. These interventions are often highly context ...

Newborn Health - World Health Organization (WHO)

Jan 1, 2012 · Perinatal asphyxia Birth asphyxia, defined as the failure to establish breathing at birth, accounts for an estimated 900,000 deaths each year and is one of the primary causes of ...

PHO Webinar: 2SLGBTQIA+ families: Unique perinatal mental ...

Apr 10, 2024 · Event Details PHO Webinar: 2SLGBTQIA+ families: Unique perinatal mental health concerns 2SLGBTQIA+ individuals face unique challenges during the perinatal period. Some ...

Sexual and Reproductive Health and Research (SRH)

Apr 25, 2023 · Maternal and perinatal health Each day, about 800 women across the world die from complications related to pregnancy or childbirth, most of which are preventable or treatable.

Maternal and Infant Health - Public Health Ontario

Jan 20, 2021 · Documents and resources related to the health of mothers and children during pregnancy and the first years of life.

WHO guide for integration of perinatal mental health in maternal ...

Sep 19, 2022 · The guide for integration of perinatal mental health in maternal and child health services outlines an evidence-informed approach describing how program managers, health service administrators and policy-makers responsible of planning and managing maternal and child health services can develop and sustain high-quality, integrated mental health ...

Evidence Brief: Exploring interventions to address perinatal ...

The time from pregnancy to childbirth is commonly referred to as the perinatal period. There are multiple working definitions for 'perinatal period'; however, for the purpose of this Evidence Brief 'perinatal mood disorders' refer to affective disorders (e.g., depression and anxiety) experienced among women during the 'perinatal period', spanning from "conception to one year ...

IPC in Perinatology_ENGLISH_2015_Revision - Public Health ...

How to cite this document: Ontario Agency for Health Protection and Promotion (Public Health Ontario), Provincial Infectious Diseases Advisory Committee. Best practices for infection prevention and control in perinatology, in all health care settings that provide obstetrical and newborn care. 1st revision, February 2015. Toronto, ON: Queen's Printer for Ontario; 2015.

Maternal and perinatal death surveillance and response

Maternal and perinatal death surveillance and response (MPDSR) is defined as an essential quality improvement intervention which permits the identification, notification, quantification and determination of causes and avoidability of maternal and neonatal deaths and stillbirth with the goal of orienting the measures necessary for their prevention.

Newborn mortality - World Health Organization (WHO)

Mar 14, 2024 · WHO fact sheet on newborn mortality, including key facts, causes, priority strategies, newborn care and WHO response.

Research in maternal, perinatal, and adolescent health

Mar 30, 2019 · WHO develops guidelines on a broad array of clinical, public health, health system, health promotion and implementation strategies. These interventions are often highly context-specific, with multiple factors that directly and indirectly impact the health and societal outcomes, and as part of these strategies, to identify key research priorities that could accelerate ...

Newborn Health - World Health Organization (WHO)

Jan 1, 2012 · Perinatal asphyxiaBirth asphyxia, defined as the failure to establish breathing at birth, accounts for an estimated 900,000 deaths each year and is one of the primary causes of early neonatal mortality. The most common cause of perinatal asphyxia is complications during childbirth. Guidelines for neonatal resuscitation emphasise the importance of drying, ...

PHO Webinar: 2SLGBTQIA+ families: Unique perinatal mental ...

Apr 10, 2024 · Event Details PHO Webinar: 2SLGBTQIA+ families: Unique perinatal mental health concerns 2SLGBTQIA+ individuals face unique challenges during the perinatal period. Some of these challenges may be related to pre-existing or new mental health issues, partners and relationship dynamics, or parenting contexts that might be different from those of cisgender ...

Enhance your skills with our comprehensive perinatal mood disorders training. Discover how to support new parents effectively. Learn more today!

[Back to Home](#)